

UPRF 97-01
January 1997

Spin Networks and Recoupling in Loop Quantum Gravity^{1,2}

Roberto De Pietri

Dipartimento di Fisica,
Università degli studi di Parma
and
INFN, Gruppo Collegato di Parma,
ITALY.

Abstract

I discuss the role played by the spin-network basis and recoupling theory (in its graphical tangle-theoretic formulation) and their use for performing explicit calculations in loop quantum gravity. In particular, I show that recoupling theory allows the derivation of explicit expressions for the eigenvalues of the quantum volume operator. An important side result of these computations is the determination of a scalar product with respect to which area and volume operators are symmetric, and the spin network states are orthonormal.

¹To appear in the Proceedings of the 2nd Conference on Constrained Dynamics and Quantum Gravity, Santa Margherita, Italy, 17-21 September 1996.

² This work has been partially supported by the INFN grant “Iniziativa specifica FI-41” (Italy).

1 Introduction

A promising attempt towards a quantum theory of gravitation is provided by the loop quantization [1]. It amounts to the direct canonical quantization of the (Poisson bracket) holonomy algebra generated by:

$$\mathcal{T}[\alpha] = -\text{Tr}[U_\alpha], \quad (1)$$

$$\mathcal{T}^a[\alpha](s) = -\text{Tr}[U_\alpha(s, s)\tilde{E}^a(s)], \quad (2)$$

where U_α is the parallel propagator of the Ashtekar-Sen connection A_a along a loop α . The crucial point is that to each loop α is uniquely associated a $\mathcal{T}$ observable. This suggests to use as carrier space of the representation a suitable subspace $\mathcal{V}$ of the free loop algebra defined as the finite linear combinations Φ of finite products of loops, as in:

$$\Phi = c_0 + \sum_i c_i[\alpha_i] + \sum_{jk} c_{jk}[\alpha_j][\alpha_k] + \dots \quad (3)$$

In this talk I shall summarize the results obtained in [5]; in particular, I shall show that the $\mathcal{V}$ -space admits a tangle-theoretic description in terms of the Temperley-Lieb recoupling theory [6], and that a convenient basis in this space is given by the spin-network states.

It is important to note that an element of the $\mathcal{T}$ holonomy algebra

$$\mathcal{T}[\Phi] = c_0 + \sum_i c_i \mathcal{T}[\alpha_i] + \sum_{ij} c_{ij} \mathcal{T}[\alpha_i] \mathcal{T}[\alpha_j] + \dots$$

is associated to each element of the free loop algebra, and that the connection representation is defined by introducing an Hilbert space structure on a suitable extensions $\overline{\mathcal{A}/\mathcal{G}}$ of this $\mathcal{T}$ algebra [2]. The relation between the two representations is given by the loop transform [3]. For a more detailed analysis of this point see [4].

2 The Carrier Space of the Loop Representation and the Spin-Networks Basis

The loop representation of quantum gravity is assumed to be the linear representation of the Poisson algebra of the $\mathcal{T}$ variables over $\mathcal{V}$ defined by: $\langle \Phi | \mathcal{T}[\beta] = \langle \Phi \cdot [\beta] |$. It may happen that two elements Φ and Ψ of the free loop algebra correspond to the same function on the holonomy algebra $\mathcal{T}[\Phi] = \mathcal{T}[\Psi]$ (i.e., they give the same value for any values of the connection). As a consequence, the carrier space $\mathcal{V}$ of the representation must be defined

as the space of the equivalence classes of the free loop algebra under the equivalence defined by all the relations (Mandelstam relations)

$$\Phi \sim \Psi \quad \text{if} \quad \mathcal{T}[\Phi] = \mathcal{T}[\Psi], \quad (4)$$

namely by the equality of the corresponding holonomies. The principal consequences of the Mandelstam relations are that the $\mathcal{V}$ -space does not depend on the orientation and parameterization of the loops. Moreover, the following identities hold true: **[Retracing]**: if γ is a *segment* starting in a point of α then $[\alpha \circ \gamma \circ \gamma^{-1}] \sim [\alpha]$; **[Binor identity]**: $[\alpha] \cdot [\beta] \sim -[\alpha \#_s \beta] - [\alpha \#_s \beta^{-1}]$.

According to [5], a natural way to represent an element Φ of $\mathcal{V}$ is the following: first, one introduces the extended planar graph Γ_{ex} of Φ as the two-dimensional surface obtained by “thickening out” a planar representation of the image γ of all the loop in Φ ; then, one draws the loops of Φ over Γ_{ex} . In this way, to each element of the $\mathcal{V}$ -space corresponds an element in the algebra of all possible tangles over Γ_{ex} . Moreover, in the tangle-theoretic planar representation of the states $\langle \Phi |$, the *retracing* and *binor* identities become the following tangle identities

$$\bigcirc = -2 \quad (5)$$

$$\times = - \mid \mid - \smile \quad . \quad (6)$$

These identities are the key axioms of the tangle-theoretic recoupling theory [6], if the value of the A parameter set to -1 . This gives the possibility of using the whole machinery of the tangle-theoretic recoupling theory [6] for dealing with the graphical representation of the states. In particular, a basis in this tangle-theoretic algebra is provided by the spin-networks, where a spin network S is the sum of tangles given as follow: consider a three-valent “virtual” graph Γ^v over Γ_{ex} and a “compatible coloring” $\{p_e\}$ of the edges of Γ^v ; replace each edge with the full-antisymmetric sum of p_e parallel lines; then, in each three-valent vertices connect all the incoming lines in the unique possible planar way.

In this way, any element of the $\mathcal{V}$ -space can be characterized in terms of the tangle-theoretic **spin network quantum state** $\langle S | = (\gamma, \Gamma_{ex}, S)$, defined as the element of $\mathcal{V}$ determined by the graph γ , its extended planar graph Γ_{ex} , and the spin-network tangle S over Γ_{ex} . These states are essentially characterized by the graph γ , the number p_e of loops in each edge e of γ , and the recoupling (in terms of three-valent vertices) of the edges connected to each vertex of γ (see figure 1).

Note that, representing a vertex of a spin network state $\langle S |$ by means of




Figure 1: An example of spin-network state and of its extended planar representation

the portion of its spin-network S contained in the vertex

$$\left\langle V^{(n)}[P_0, \dots, P_{n-1}](K_{I_i}) \right| = \left\langle \begin{array}{c} P_2 \quad \dots \quad P_{n-3} \\ \diagdown \quad \diagup \quad \diagdown \quad \diagup \\ P_1 \quad i_2 \quad i_3 \quad \dots \quad i_{n-2} \\ \diagup \quad \diagdown \quad \diagup \quad \diagdown \\ P_0 \quad i_1 \quad i_{n-1} \end{array} \begin{array}{c} P_{n-2} \\ \diagup \quad \diagdown \\ P_{n-1} \end{array} \right|, \quad (7)$$

and considering two vertices, $V_n[a_1, \dots, a_n]$ and $V'_n[b_1, \dots, b_n]$, they can be connected only if all the corresponding incoming lines have the same colors. In this case, it is possible to attach the a_i lines and the corresponding b_i lines to each other. Then, it is obtained a tangle over Γ_{ex} constituted only of loops contractible to a point. By recoupling, a tangle in which all the loops are contractible to a point reduce to a number. The operation of computing this number is named chromatic evaluation. This evaluation, denoted by $\langle V_n(a_1, \dots, a_n) | V'_n(b_1, \dots, b_n) \rangle$, is a scalar product in the space of the possible vertices. In particular, the chromatic evaluation of a 2-vertex (i.e. of a line) and that of a 3-vertex will be denoted as Δ function and θ function, respectively:

$$\Delta_n = \langle V_2[n], V_2[n] \rangle = \text{diagram of a line with a loop labeled } n \quad (8)$$

$$\theta(a, b, c) = \langle V_3[a, b, c], V_3[a, b, c] \rangle = \text{diagram of a triangle with lines labeled } a, b, c \quad (9)$$

For the explicit values of this evaluation and all the details see [5] and [6]. In this tangle-theoretic interpretation of the loop representation, one still has the analogous of the Wigner-Eckart theorem: the recoupling theorem of [6] (pg. 60) states, as a tangle relation, that

$$\begin{array}{c} b \quad c \\ \diagdown \quad \diagup \\ \bullet \quad \bullet \\ \diagup \quad \diagdown \\ a \quad d \end{array} = \sum_i \left\{ \begin{array}{ccc} a & b & i \\ c & d & j \end{array} \right\} \begin{array}{c} b \quad c \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ a \quad d \end{array}, \quad (10)$$

where the quantities $\left\{ \begin{smallmatrix} a & b & i \\ c & d & j \end{smallmatrix} \right\}$ are $su(2)$ six-j symbols (normalized as in [6]).

One of the main advantages of the spin-network basis is that the action of the $\mathcal{T}^a[\alpha](s)$ operators is particularly simple. In fact, one has:

$$= p \Delta^a[\beta, s] \quad (11)$$

where $\Delta^a[\beta, s] = \int_{\beta} d\lambda \dot{\beta}^a(\lambda) \delta^3[\beta(\lambda), s]$. Indeed the action of the operators corresponds to adding a new vertex of the kind shown in eq. (11) at the intersection point s between α and β . This gives the possibility of expressing the action of any operator on the spin network basis as a product of three parts: an analytic regularization factor, a graphical part (due the a possible change of γ), and a vertex part (see [8]). The vertex part, connected with the reduction to elementary vertices, can be computed using the recoupling theory and the associated chromatic evaluations.

3 The Volume Operator

As an example of application of the recoupling theory, I show how is possible to obtain explicit formulas for the eigenvalues of the volume operator. In [5], it has been shown that the action of the volume operator is given by a regularization factor and a vertex part. The following final expression for the action of the volume operator on a given spin-network has been obtained

$$\hat{V}[\mathcal{V}] = \sum_{i \in \{S \cap \mathcal{V}\}} \hat{V}_i, \quad (12)$$

$$\hat{V}_i = l_0^3 \sqrt{\sum_{r \neq s \neq t=0}^{n_i-1} \left| \frac{i}{16 \cdot 3!} \hat{W}_{[rts]}^{(n_i)} \right|}.$$

The first sum is over all the vertices and the second sum is over the triples of edges adjacent to each vertex. In [5], it has also been shown that the vertex operators $\hat{W}_{[rts]}^{(n_i)}$ are represented by diagonalizable matrices with real eigenvalues. In fact, a normalization of the vertex exists such that, in this basis, the vertex operator $\hat{W}_{[rts]}^{(n_i)}$ are represented by the real and antisymmetric matrices

$$\left\langle V^{(n_i)}(K_{I_i}) \right|_N \hat{W}_{[rts]}^{(n_i)} = \sum_{\bar{K}_{I_i}} \hat{W}_{[rts] K_{I_i}}^{(n_i) \bar{K}_{I_i}} \left\langle V^{(n_i)}(\bar{K}_{I_i}) \right|_N$$

In the case of 4-valent vertices, this normalization is explicitly given by:

$$\langle V^{(4)}(i) \rangle_N = \sqrt{\frac{\text{Diagram 1}}{\text{Diagram 2}}} \left| \text{Diagram 3} \right|, \quad (13)$$

Diagram 1: A square with a vertical line through the center, labeled i at the top. Diagram 2: Two circles, the left one labeled $\frac{a}{b}$ and $\frac{b}{i}$, the right one labeled $\frac{c}{d}$ and $\frac{d}{i}$. Diagram 3: A vertex labeled i with four edges labeled a, b, c, d .

and the matrix elements of the vertex operator are given by the chromatic evaluation:

$$\tilde{W}_{[012]}^{(4)}(a, b, c, d)_i^k = a \cdot b \cdot c \cdot \sqrt{\frac{\text{Diagram 1}}{\text{Diagram 2}}} \cdot \text{Diagram 3}. \quad (14)$$

Diagram 1: A square with a vertical line through the center, labeled i at the top and k at the bottom. Diagram 2: Four circles, the first two labeled $\frac{a}{b}$ and $\frac{a}{k}$, the last two labeled $\frac{c}{d}$ and $\frac{c}{k}$. Diagram 3: A complex diagram with vertices labeled a, b, c, d, i, k and edges labeled 2 .

The matrices $\tilde{W}_{[012]}^{(4)}(a, b, c, d)_i^k$ have elements different from zero only if $|i - k| = 2$. Using the equations

$$\begin{array}{c} k \\ \bullet \\ a \bullet \\ i \bullet \end{array} \begin{array}{c} \bullet \\ \bullet \\ \bullet \end{array} \begin{array}{c} c \\ \bullet \\ \bar{c} \end{array} d = \frac{\text{Diagram 1}}{\text{Diagram 2}} \cdot \begin{array}{c} k \\ \bullet \\ a \bullet \\ i \bullet \end{array} \begin{array}{c} \bullet \\ \bullet \\ \bullet \end{array} \quad (15)$$

Diagram 1: A square with vertices labeled k, a, c, i and edges labeled $a, c, \bar{c}, i$. Diagram 2: A circle labeled $\frac{k}{a}$ and $\frac{a}{i}$.

$$r \cdot \begin{array}{c} p \\ \bullet \\ 2 \\ \bullet \\ r \end{array} \begin{array}{c} q \\ \bullet \\ \bullet \end{array} = p \cdot \begin{array}{c} p \\ \bullet \\ 2 \\ \bullet \\ r \end{array} \begin{array}{c} q \\ \bullet \\ \bullet \end{array} + q \cdot \begin{array}{c} p \\ \bullet \\ 2 \\ \bullet \\ r \end{array} \begin{array}{c} q \\ \bullet \\ \bullet \end{array}, \quad (16)$$

it is possible to reduce eq. (14) to the following chromatic evaluation:

$$\tilde{W}_{[012]}^{(4)}(a, b, c, d)_i^k = \begin{array}{c} \frac{k}{2} \\ \bullet \\ \frac{2}{i} \end{array} \cdot \sqrt{\frac{\text{Diagram 1}}{\text{Diagram 2}}} \cdot \left[b \frac{\text{Diagram 3}}{\frac{b}{2}} - k \frac{\text{Diagram 4}}{\frac{k}{2}} \right] \cdot \left[b \frac{\text{Diagram 5}}{\frac{k}{2}} \right] \cdot \left[c \frac{\text{Diagram 6}}{\frac{k}{2}} \right]. \quad (17)$$

Diagram 1: A square with a vertical line through the center, labeled i at the top and k at the bottom. Diagram 2: Four circles, the first two labeled $\frac{a}{b}$ and $\frac{a}{k}$, the last two labeled $\frac{c}{d}$ and $\frac{c}{k}$. Diagram 3: A square with vertices labeled $2, 2, b, b$ and edges labeled $2, 2, b, b$. Diagram 4: A square with vertices labeled $2, 2, k, i$ and edges labeled $2, 2, k, i$. Diagram 5: A square with vertices labeled $k, 2, b, b$ and edges labeled $k, 2, b, b$. Diagram 6: A square with vertices labeled $k, 2, c, c$ and edges labeled $k, 2, c, c$.

Define $t = (i+k)/2$ and $\epsilon = (k-i)/2$. The matrix element $\tilde{W}_{[012]}^{(4)}(a, b, c, d)_i^k(a, b, c, d)$ is different from zero only if $\epsilon = \pm 1$ and all the 3-vertices in equation (17) are admissible. Indeed, by performing the chromatic evaluations of eq. (17),

one gets the following explicit formula for the matrix elements ($\epsilon = \pm 1$):

$$\begin{aligned} \tilde{W}_{[012]}^{(4)}(a, b, c, d)_{t-\epsilon}^{t+\epsilon} &= -\epsilon(-1)^{\frac{a+b+c+d}{2}} \\ &\left[\frac{1}{4t(t+2)} \frac{a+b+t+3}{2} \frac{c+d+t+3}{2} \right. \\ &\frac{1+a+b-t}{2} \frac{1+a+t-b}{2} \frac{1+b+t-a}{2} \\ &\left. \frac{1+c+d-t}{2} \frac{1+c+t-d}{2} \frac{1+d+t-c}{2} \right]^{\frac{1}{2}} \end{aligned} \quad (18)$$

This formula can be used to obtain explicit formulas for the eigenvalues of the volume operator. For example, in the case $d = a + b + c - 2$, where the matrices $\tilde{W}_{[rst]}$ are two-dimensional, one obtains the following result for the (degenerate) eigenvalue of the volume associated to the 4-vertex $\langle V^{(4)}[a, b, c, a + b + c - 2] \rangle$:

$$v(a, b, c, d) = \frac{l_0^3}{\sqrt{2}} \left[\frac{a b c (a + b + c)}{16} \right]^{\frac{1}{4}}, \quad (19)$$

that can be compared with the results of [9].

4 The Normalized State and the Scalar Product

A scalar product between two spin-network states can be defined by assuming that it is different from zero only if the two states have exactly the same graph and the same coloring of the real edges. Then, the normalization (13), with respect to which the volume operator is represented by symmetrical matrices, determines this scalar product uniquely. Its value is determined by the chromatic evaluation of the vertices:

$$\langle s, s' \rangle = \delta_{\gamma, \gamma'} \prod_{e \in \mathcal{E}_s} \frac{\delta_{n_e, n'_e}}{\Delta_{n_e}} \prod_{i \in \mathcal{V}_s} \langle V_i, V'_i \rangle, \quad (20)$$

where the products are extended only to the real edge and to the real vertices, and $\langle V_i, V'_i \rangle$ is the chromatic evaluation obtained by gluing the vertices V_i and V'_i . In [4], it has been shown that the scalar product defined in this way is precisely the loop transform [3] of the Ashtekar-Lewandowski measure [2].

I thank warmly Massimo Pauri for a critical reading of the manuscript.

References

- [1] C. Rovelli and L. Smolin *Phys. Rev. Lett.* **61**(1988)1155; *Nucl. Phys.* **B331**(1990)80. For a recent review see C. Rovelli, gr-qc/9606090 and reference therein.
- [2] A. Ashtekar, J. Lewandowski, D. Marolf, J. Mourão and T. Thiemann, *J. Math. Phys.* **36**(1995),6456, and reference therein.
- [3] A. Ashtekar A and C. Isham, *Class. Quantum Grav.* **9**(1992)1433.
- [4] R. De Pietri, *Class. Quantum Grav.* **14**(1997)53.
- [5] R.DePietri and C. Rovelli, *Phys. Rev. D* **54**(1996)2664.
- [6] L.H. Kauffman and L.S. Lins *Temperley-Lieb Recoupling Theory and Invariants of 3-Manifolds* (Princeton University Press, Princeton (NJ), 1994).
- [7] C. Rovelli and L. Smolin *Phys. Rev. D* **53**(1995)5743; *Nucl. Phys.* **B442**(1995)593.
- [8] R. Borissov, in this volume.
- [9] J. Lewandowski, *Class. Quantum Grav.* **14**(1997)71. T. Thiemann, gr-qc/9606091.